

UNIVERSIDAD AUTÓNOMA METROPOLITANA - AZCAPOTZALCO
TRIMESTRE: INVIERNO DE 2020.

CÁLCULO DIFERENCIAL
EXAMEN # 1 (FORMA REMOTA). - AA
FECHA: SÁBADO 6 DE JUNIO DE 2020: 14:30 HORAS

Nombre:

ANSWER KEY

Instrucciones:

- El examen consta de SEIS problemas con diferentes puntajes.
- Tienen **una hora con treinta (30) minutos** para resolverlos.
- El examen es **INDIVIDUAL**. Está prohibido recibir ayuda de terceras personas o usar recursos no especificados.
- Pueden usar sus libros, apuntes y calculadora sencilla. Cite cuando use libros o apuntes, o su calculadora.
- Para recibir puntaje: Conteste correctamente. Escriba de forma clara y concisa. Entregue su trabajo limpio y con sus ideas en orden. **SIMPLIFIQUE** y muestre todas sus cuentas. **EXPLIQUE, ARGUMENTE y JUSTIFIQUE** sus respuestas.
- Problema **SIN explicación, desarrollo, justificación o argumento** vale **CERO** puntos.

PROBLEMAS

- (0) No olvide elaborar la carátula del examen y anexarla en su escaneado.
- (1) (10 puntos.) Calcule la ecuación de la recta tangente a la gráfica de la función $f(x) = -\frac{1}{(17-x)^{3/2}}$ en el punto $(1, -1/4)$. Use la definición de derivada.
- (2) Calcule:
(a) (10 puntos) la derivada de $f(x) = \sin(\tan^2(\cos 3x))$.
(b) (10 puntos) $(\sin x)^{(139)}$.
- (3) (10 puntos.) Encuentre la ecuación de la recta ortogonal a la curva $x^2 + y^2 = 16$ en el punto $(2, 2)$.
- (4) (20 puntos.) Un carro se mueve a lo largo de una autopista y su posición está dada por la función $x(t) = t^4 - 4t^2$. ¿En qué instantes el carro frena?
- (5) (20 puntos.) Imagine una playa totalmente recta. Un barco va navegando de forma paralela a la playa durante la noche y un faro lo va siguiendo. El barco navega a una distancia de 500 metros de la playa a una velocidad de 200 metros por minuto. El rayo del faro hacia el barco hace un ángulo con la playa. Cuando el barco está a 1000 metros del faro, ¿a qué velocidad angular se mueve el faro?
- (6) (20 puntos.) En clase se vio cuáles son las diferencias y las similitudes cuando se quieren calcular la recta tangente a una curva y la velocidad instantánea de una partícula. ¿Cuáles son? Describalas y explique.
- (*) **FÓRMULAS.**
(a) $\sin^2 \alpha + \cos^2 \alpha = 1$.
(b) $\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$
(c) $\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$
(d) $\cos^2 \alpha = \frac{1 + \cos 2\alpha}{2}$.
(e) $\sin^2 \alpha = \frac{1 - \cos 2\alpha}{2}$.

Examen # 1ANSWER KEY

① We have the function $f(x) = -\frac{1}{\sqrt{17-x}}$ and $f(1) = -\frac{1}{4}$, so that the graph passes through $(1, -1/4)$. If we want to find the slope of the tangent line, using the derivative function: $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$. Using this definition:

$$f'(x) = \lim_{h \rightarrow 0} \frac{\left(-\frac{1}{\sqrt{17-(x+h)}}\right) - \left(-\frac{1}{\sqrt{17-x}}\right)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\left(\frac{-\sqrt{17-x} + \sqrt{17-(x+h)}}{\sqrt{17-(x+h)} \sqrt{17-x}}\right)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{1}{h} \frac{\sqrt{17-(x+h)} - \sqrt{17-x}}{\sqrt{17-(x+h)} \sqrt{17-x}} \frac{(\sqrt{17-(x+h)} + \sqrt{17-x})}{(\sqrt{17-(x+h)} + \sqrt{17-x})}$$

$$= \lim_{h \rightarrow 0} \frac{1}{h} \frac{(17-(x+h)) - (17-x)}{\sqrt{17-(x+h)} \sqrt{17-x} (\sqrt{17-(x+h)} + \sqrt{17-x})}$$

$$= \lim_{h \rightarrow 0} \frac{1}{h} \frac{-h}{\sqrt{17-(x+h)} \sqrt{17-x} (\sqrt{17-(x+h)} + \sqrt{17-x})}$$

$$= \lim_{h \rightarrow 0} \frac{-1}{(\sqrt{17-(x+h)} \sqrt{17-x} (\sqrt{17-(x+h)} + \sqrt{17-x}))}$$

Now, we can directly substitute $h=0$; to evaluate limit:

$$= \frac{-1}{\sqrt{17-x} \sqrt{17-x} (\sqrt{17-x} + \sqrt{17-x})}$$

$$= \frac{-1}{(17-x) 2\sqrt{17-x}}$$

$$= -\frac{1}{2(17-x)\sqrt{17-x}}$$

Too, the derivative function is:

$$f'(x) = \frac{-1}{2(17-x)^{3/2}}$$

At the value $x=1$, we get the slope of the tangent line:

$$m = f'(1) = \frac{-1}{2(16)^{3/2}} = \frac{-1}{2(4^3)} = \frac{-1}{2^7}$$

is. $m = \frac{1}{128}$. And the equation of the tangent

line in point-slope form is

$$y - \left(-\frac{1}{4}\right) = \frac{-1}{128}(x-1)$$

is.
$$y + \frac{1}{4} = \frac{-1}{128}(x-1)$$

In slope-y-intercept form:

$$y = \frac{-1}{128}x + \frac{1}{128} - \frac{1}{4} = \frac{-1}{128}x + \frac{1-32}{128}$$

$$y = \frac{-1}{128}x - \frac{31}{128}$$

(2) (a) We use the Chain rule to compute:

$$\begin{aligned} \frac{d}{dx} f(x) &= \frac{d}{dx} \left(\sin(\tan^2(\cos 3x)) \right) \\ &= \cos(\tan^2(\cos 3x)) \frac{d}{dx} (\tan^2(\cos 3x)) \quad \checkmark \text{Chain rule} \\ &= \cos(\tan^2(\cos 3x)) \cdot 2 \tan(\cos 3x) \frac{d}{dx} \tan(\cos 3x) \quad \checkmark \text{Chain rule} \\ &= \cos(\tan^2(\cos 3x)) \cdot 2 \tan(\cos 3x) \sec^2(\cos 3x) \cdot \frac{d}{dx} (\cos 3x) \quad \checkmark \text{Chain rule} \\ &= \cos(\tan^2(\cos 3x)) \cdot 2 \tan(\cos 3x) \sec^2(\cos 3x) (-\sin 3x) \frac{d}{dx} (3x) \quad \checkmark \text{Chain rule} \\ &= \cos(\tan^2(\cos 3x)) \cdot 2 \tan(\cos 3x) \sec^2(\cos 3x) (-1) \sin(3x) \cdot 3. \end{aligned}$$

Rearranging terms.

$$\begin{aligned} \frac{df}{dx} &= \frac{d}{dx} \left(\sin(\tan^2(\cos 3x)) \right) = \\ &= -6 \cos(\tan^2(\cos 3x)) \tan(\cos 3x) \sec^2(\cos 3x) \sin(3x) \end{aligned}$$

(b) Compute $(\sin x)^{(139)} = \frac{d^{139}}{dx^{139}} (\sin x)$.

Notice that 136 is a multiple of 4. Thus.

$$\frac{d^{136}}{dx^{136}} \sin x = \sin x.$$

$$\text{Hence } \frac{d^{139}}{dx^{139}} (\sin x) = \frac{d^3}{dx^3} \left(\frac{d^{136}}{dx^{136}} \sin x \right) = \frac{d^3}{dx^3} (\sin x) =$$

$$= \frac{d^2}{dx^2} (\cos x) = \frac{d}{dx} (-\sin x) = -\cos x$$

$$\Rightarrow \boxed{(\sin x)^{(139)} = -\cos x}$$

③ We have the function $y = f(x)$ implicitly defined by:

$$x^3 + y^3 = 16.$$

We observe that $(2, 2)$ belongs to the curve since

$$2^3 + 2^3 = 8 + 8 = 16 \quad \checkmark$$

Now, compute the derivative on each side of eq'n:

$$3x^2 + 3y^2 y' = 0, \text{ by chain rule.}$$

$$\text{Then, } y' = -\frac{x^2}{y^2}.$$

$$\text{At } (2, 2), m_{\text{tan}} = y' \Big|_{(2,2)} = -\frac{2^2}{2^2} = -1,$$

this is the slope of the tangent line.

The slope of the orthogonal line is:

$$m = -\frac{1}{m_{\text{tan}}} = -\frac{1}{-1} = 1$$

Therefore, the equation of the orthogonal line

is:

$$y - 2 = 1 \cdot (x - 2)$$

$$\text{ie } y - 2 = x - 2$$

$$\Rightarrow \boxed{y = x}$$

④ The trajectory of the particle is:

$$x(t) = t^4 - 4t^3$$

The derivative is the velocity.

$$\dot{x}(t) = 4t^3 - 12t^2.$$

The acceleration is the second derivative:


$$\ddot{x}(t) = 12t^2 - 24t.$$

We are asked ^{for} the intervals of time when the particle slows down, i.e., when \dot{x} and \ddot{x} have opposite signs.

Observe that:

$$\dot{x} = 4t^2(t - 3)$$


$$\ddot{x} = 12t(t - 2).$$



We can plot these two functions and observe when they have opposite signs.

We have:
 $v < 0, a > 0$, on $(-\infty, 0)$.

also:
 $v < 0, a > 0$, on $(2, 3)$




Then, the particle slows down if

$$t \in (-\infty, 0) \cup (2, 3)$$

(In the interval $(0, 2) \cup (3, \infty)$, v and a have same signs: speeds up).

5) Let's do a picture

Here $l = 500\text{m}$
 $v = 200\text{m/min}$



This are the only constraints.

The variables (dependent variables) are $\left. \begin{array}{l} l(t) \\ \theta(t) \end{array} \right\} x(t) = \text{projective of ship on beach.}$

$l = l(t) = \text{distance from lighthouse to ship}$

$\theta = \theta(t) = \text{angle of the ray light and the beach}$

(time is the independent variable).

Observe that:

$$\tan \theta(t) = \frac{l}{x(t)}$$

since we have a right triangle. Computing the

derivatives;
 by chain rule: $(\sec^2 \theta) \dot{\theta} = -\frac{l}{x^2} \dot{x}$

i.e. $\dot{\theta} = -\frac{l}{x^2} \dot{x} \cos^2 \theta$

But $\cos^2 \theta = \left(\frac{x}{l}\right)^2$ Then $\dot{\theta} = -\frac{l}{x^2} \dot{x} \left(\frac{x}{l}\right)^2$

i.e. $\dot{\theta} = -\frac{l}{l^2} \dot{x}$ since $\dot{x} = v = 200\text{m/sec}$

and since we want $\dot{\theta}$ when $l = 1000\text{m}$:

$$\dot{\theta} = -\frac{1}{10} \frac{1}{\text{sec}}$$

$$\dot{\theta} = -\frac{(500\text{m})(200\text{m/sec})}{(1000\text{m})^2} \Rightarrow \dot{\theta} = \frac{-100,000 \text{ m}^2/\text{sec}}{1,000,000 \text{ m}^2}$$

⑥ If we have the equation of the graph of the function $f(x)$, the graph has equation:

$$y = f(x).$$

And for $x = x_0$, then $y_0 = f(x_0)$, and the graph

passes through: $(x_0, y_0) = (x_0, f(x_0))$.

To compute the slope, we use the derivative: at $x = x_0$

$$m = f'(x_0) = \lim_{h \rightarrow 0} \frac{f(x_0+h) - f(x_0)}{h}.$$

Now, if a particle has a position x , given by the function $f(t)$, then $x = f(t)$, and the derivative at a time $t = t_0$ is the instantaneous velocity

$$v(t) = f'(t_0) = \lim_{\Delta t \rightarrow 0} \frac{f(t_0 + \Delta t) - f(t_0)}{\Delta t}.$$

We observe that it is exactly the same operation!
So, this is the similitude. There is no difference
in the mathematics, we have the same operation!

The difference is the interpretation:

For the slope, the secant line has a slope

$$\frac{f(x_0+h) - f(x_0)}{h} = \frac{\Delta y}{\Delta x},$$

while we interpret the fraction:

$$\frac{f(t_0 + \Delta t) - f(t_0)}{\Delta t} = \frac{\Delta x}{\Delta t} \text{ as an } \underline{\text{average velocity}}$$